

---

# Safety Assessment of Wheat-Derived Ingredients as Used in Cosmetics

---

Status: Tentative Report for Public Comment  
Release Date: June 19, 2020  
Panel Meeting Date: September 14-15, 2020

*All interested persons are provided 60 days from the above release date (i.e., August 18, 2020) to comment on this safety assessment and to identify additional published data that should be included or provide unpublished data which can be made public and included. Information may be submitted without identifying the source or the trade name of the cosmetic product containing the ingredient. All unpublished data submitted to CIR will be discussed in open meetings, will be available at the CIR office for review by any interested party and may be cited in a peer-reviewed scientific journal. Please submit data, comments, or requests to the CIR Executive Director, Dr. Bart Heldreth.*

The Expert Panel for Cosmetic Ingredient Safety members are: Chair, Wilma F. Bergfeld, M.D., F.A.C.P.; Donald V. Belsito, M.D.; Curtis D. Klaassen, Ph.D.; Daniel C. Liebler, Ph.D.; James G. Marks, Jr., M.D.; Lisa A. Peterson, Ph.D.; Ronald C. Shank, Ph.D.; Thomas J. Slaga, Ph.D.; and Paul W. Snyder, D.V.M., Ph.D. The Cosmetic Ingredient Review (CIR) Executive Director is Bart Heldreth, Ph.D. This safety assessment was prepared by Christina L. Burnett, Senior Scientific Analyst/Writer, CIR.

## ABSTRACT

The Expert Panel for Cosmetic Ingredient Safety (Panel) assessed the safety of 27 wheat-derived ingredients. Most of these ingredients are reported to function as skin conditioning agents in cosmetic products. Industry should continue to use good manufacturing practices to limit impurities that could be present in botanical ingredients. The Panel reviewed the available data to determine the safety of these ingredients, noting the lack of method of manufacture data for all ingredients and dermal irritation and sensitization data for one ingredient, and concluded that the available data are insufficient to make a determination of safety for these wheat-derived ingredients under the intended conditions of use in cosmetic formulations.

## INTRODUCTION

Most of wheat-derived ingredients detailed in this safety assessment are reported to function in cosmetics as skin conditioning agents, while some are reported to have other functions, such as abrasives, absorbents, antioxidants, bulking agents, film formers, flavoring agent, hair conditioning agents, and viscosity increasing agents, according to the web-based *International Cosmetic Ingredient Dictionary and Handbook* (wINCI; *Dictionary*; see Table 1).<sup>1</sup> Functions such as skin bleaching agent (for *Triticum Vulgare* (Wheat) Germ Extract) are not considered cosmetic functions in the United States (US) and, therefore, are not addressed in this assessment. This assessment of the safety of the following 27 wheat-derived ingredients is based on the data contained in this report:

<i>Triticum Aestivum</i> (Wheat) Flour Lipids	<i>Triticum Vulgare</i> (Wheat) Flour Lipids
<i>Triticum Aestivum</i> (Wheat) Germ Extract	<i>Triticum Vulgare</i> (Wheat) Germ
<i>Triticum Aestivum</i> (Wheat) Leaf Extract	<i>Triticum Vulgare</i> (Wheat) Germ Extract
<i>Triticum Aestivum</i> (Wheat) Peptide	<i>Triticum Vulgare</i> (Wheat) Germ Powder
<i>Triticum Aestivum</i> (Wheat) Seed Extract	<i>Triticum Vulgare</i> (Wheat) Germ Protein
<i>Triticum Monococcum</i> (Wheat) Seed Extract	<i>Triticum Vulgare</i> (Wheat) Gluten*
<i>Triticum Monococcum</i> (Wheat) Stem Water	<i>Triticum Vulgare</i> (Wheat) Gluten Extract
<i>Triticum Spelta</i> Seed Water	<i>Triticum Vulgare</i> (Wheat) Kernel Flour*
<i>Triticum Turgidum Durum</i> (Wheat) Seed Extract	<i>Triticum Vulgare</i> (Wheat) Protein
<i>Triticum Vulgare/Aestivum</i> (Wheat) Grain Extract	<i>Triticum Vulgare</i> (Wheat) Seed Extract
<i>Triticum Vulgare</i> (Wheat) Bran	<i>Triticum Vulgare</i> (Wheat) Sprout Extract
<i>Triticum Vulgare</i> (Wheat) Bran Extract	<i>Triticum Vulgare</i> (Wheat) Straw Water
<i>Triticum Vulgare</i> (Wheat) Bran Lipids	Wheat Germ Glycerides*
<i>Triticum Vulgare</i> (Wheat) Flour Extract	

\*Previously reviewed ingredients

The safety of Wheat Flour, now known as *Triticum Vulgare* (Wheat) Kernel Flour, was previously reviewed by the Panel and the conclusion of “safe ... in the present practices of use and concentration” was published in 1980,<sup>2</sup> and reaffirmed in a re-review that was published in 2003.<sup>3</sup> The safety of *Triticum Vulgare* (Wheat) Gluten and Wheat Germ Glycerides was also previously reviewed by the Panel and the conclusion of “safe when incorporated in cosmetic products and constitute no risk to the public in its present cosmetic use of these products” was published in 1980,<sup>4</sup> and reaffirmed in the re-review published in 2003.<sup>3</sup> Because it has been more than 15 years since the safety of *Triticum Vulgare* (Wheat) Gluten, *Triticum Vulgare* (Wheat) Kernel Flour, and Wheat Germ Glycerides were reviewed, these ingredients are included in this safety assessment for re-review. Excerpts from the summaries of the 1980 reports are disseminated throughout the text of this document, as appropriate, and are identified by *italicized text*.

The Panel has reviewed the safety of several additional wheat-derived ingredients including Wheat Amino Acids,<sup>5</sup> *Triticum Vulgare* (Wheat) Starch,<sup>2,3,6</sup> Hydrolyzed *Triticum Spelta* Starch,<sup>6</sup> Hydrolyzed Wheat Starch,<sup>6</sup> *Triticum Aestivum* (Wheat) Germ Oil,<sup>7</sup> *Triticum Vulgare* (Wheat) Germ Oil,<sup>3,7,8</sup> *Triticum Vulgare* (Wheat) Germ Oil Unsaponifiables,<sup>7</sup> Hydrogenated Wheat Germ Oil,<sup>7</sup> and Hydrogenated Wheat Germ Oil Unsaponifiables.<sup>7</sup> The Panel concluded that all of these ingredients are safe in the present practices of use and concentration in cosmetics. The Panel also reviewed Hydrolyzed Wheat Protein and Hydrolyzed Wheat Gluten, and concluded that these ingredients are safe for use in cosmetics when formulated to restrict peptides to an average molecular weight of 3500 Daltons (Da) or less.<sup>9</sup> This conclusion is in response to reports of type 1 immediate hypersensitivity reactions that occurred in sensitized individuals following exposure to cosmetic products that contained one of these two ingredients with molecular weights greater than this limit. Polypeptides must be at least 30 amino acids long (i.e. molecular weights of about 3570 Da or more) to elicit type 1 hypersensitivity reactions (i.e. to interact with the two IgE-binding epitopes).

Most of the ingredients reviewed in this safety assessment may be consumed as food, and daily exposure from food use would result in much larger systemic exposures than those from use in cosmetic products. The primary focus of the safety assessment of these ingredients as used in cosmetics is on the potential for effects from topical exposure. Proteins from wheat in the diet, specifically wheat gluten, are associated with adverse health conditions (such as celiac disease and dermatitis herpetiformis) in a small portion of the general population. Since the maximum dose of gluten possible from cosmetics is low, there is little likelihood that cosmetic products could precipitate a flare-up of either gastrointestinal or cutaneous symptoms.<sup>4,10</sup>

This safety assessment includes relevant published and unpublished data that are available for each endpoint that is evaluated. Published data are identified by conducting an exhaustive search of the world's literature. A listing of the search engines and websites that are used and the sources that are typically explored, as well as the endpoints that the Panel typically evaluates, is provided on the Cosmetic Ingredient Review (CIR) website (<https://www.cir-safety.org/supplementaldoc/preliminary-search-engines-and-websites>; <https://www.cir-safety.org/supplementaldoc/cir-report-format-outline>). Unpublished data are provided by the cosmetics industry, as well as by other interested parties.

**Note:** In many of the published studies, it is not known how the substance being tested compares to the cosmetic ingredient. Therefore, if it is not known whether the substance being discussed is a cosmetic ingredient, the test substance will be identified as “wheat...” (e.g., wheat germ extract or wheat flour); if it is known that the substance is a cosmetic ingredient, the *Dictionary* nomenclature “*Triticum Aestivum...*” or “*Triticum Vulgare...*” (e.g., *Triticum Aestivum* (Wheat) Germ Extract or *Triticum Vulgare* (Wheat) Kernel Flour) will be used. Of additional note, the accepted scientific name for both *Triticum vulgare* and *Triticum spelta* is *Triticum aestivum*.<sup>1,11</sup>

## **CHEMISTRY**

### **Definition and Plant Identification**

The definitions of the ingredients included in this review are provided in Table 1.<sup>1</sup> *Triticum* wheat species have been used as food staples for 10,000 years and originated in the Middle East.<sup>12</sup> *Triticum aestivum* L. is the most cultivated cereal grain in the world, making up about a third of total cereal grains.<sup>13</sup> *Triticum monococcum* is also known as eikorn and is native to eastern Europe and western Asia.<sup>12</sup> *Triticum turgidum* var. durum is also known as durum wheat, hard wheat, or macaroni or pasta wheat, and is commonly cultivated in the Mediterranean region.<sup>11,12</sup> *Triticum spelta* is considered to be an ancient wheat grain that was once prominently grown in ancient Egypt and Italy and is now grown in northern and eastern European countries.<sup>12,14</sup> The accepted scientific name for both *Triticum vulgare* and *Triticum spelta* is *Triticum aestivum*.<sup>1,11</sup>

Table 2 lists the generic definitions of the parts of plants that are most pertinent to the ingredients in this report.<sup>1</sup> The wheat plant is comprised of a root and shoot system.<sup>15</sup> There are two types of roots: the seminal roots and the nodal roots (crown or adventitious roots). The shoot is made up of a series of repeating units or phytomers, each potentially having a node, a leaf, an elongated internode, and a bud in the axil of the leaf. The dry fruit of the wheat plant may be referred to as the seed, grain, or kernel. The embryo of the seed is also called the germ and is comprised of the scutellum, the plumule (shoot), and the radicle (primary root). The scutellum secretes some of the enzymes involved in germination and absorbs the soluble sugars that are breakdown products of the starch in the endosperm. The endosperm is surrounded by the metabolically active cell layer called the aleurone layer, the testa (seed coat), and the pericarp (fruit coat).

#### **Triticum Vulgare (Wheat) Germ**

Wheat germ is comprised of the embryonic axis and the scutellum of wheat seed and represents about 2.5% - 3.8% of total seed weight.<sup>16</sup> It is a by-product of the flour milling industry.

### **Physical and Chemical Properties**

#### **Triticum Aestivum (Wheat) Leaf Extract**

The pH of wheatgrass (the young leaves of wheat) is reported to be 7.4.<sup>17</sup>

#### **Triticum Vulgare (Wheat) Kernel Flour**

The pH of refined wheat flours is approximately 6.0 - 6.1, while the pH of whole wheat flour is approximately 6.2.<sup>13</sup>

### **Method of Manufacture**

#### **Triticum Vulgare (Wheat) Bran Extract**

A supplier reported that its 4 different *Triticum Vulgare* (Wheat) Bran Extract products are derived from food-grade plant material and are extracted at “considerate” temperatures during a fixed time.<sup>18</sup> The resulting material is sterile filtered at the end of fabrication. Glycerin, propylene glycol, and soybean oil were used as the solvents for these products. (Table 3)

#### **Triticum Vulgare (Wheat) Kernel Flour**

Wheat flour is produced by different forms of milling or grinding, of the grain's endosperm.<sup>13,19</sup> The different forms of compression and shear allow for producing different grain sizes for different end uses or flour types.

#### **Triticum Vulgare (Wheat) Germ**

Wheat germ is a by-product of flour milling and is produced by mechanical separation of the germ from whole wheat.<sup>16</sup>

#### **Triticum Vulgare (Wheat) Gluten**

*Triticum Vulgare* (Wheat) Gluten is prepared by water washing wheat flour then drying the insoluble matter with careful temperature control.<sup>4</sup> The cream-tan powder derived from this process is food grade.

## Wheat Germ Glycerides

Wheat Germ Glycerides are produced through the transesterification of wheat germ oil (from conventional milling processes) with glycerin.<sup>4</sup>

### **Composition/Impurities**

Yields of constituents in wheat have been found to be dependent on extraction methods and growing conditions such as soil composition, climate, duration of growth period, and height at harvest.<sup>17</sup> Additionally, different plant parts have different constituent compositions.

#### Triticum Aestivum (Wheat) Leaf Extract

At least 41 flavonoid derivatives have been identified in wheat leaf extract (extracted with 80% methanol) using liquid chromatography/mass spectrometry (LC/MS).<sup>20</sup> The identity of flavones, a subgroup of flavonoids, comprised therein were luteolin, tricetin, apigenin, chrysoeriol, and tricrin.

#### Triticum Vulgare/Aestivum (Wheat) Grain Extract

Total phenolic content of wheat grain extract (as 80% methanol extracts) was 5.1 - 6.8  $\mu\text{g}$  ( $\pm$ )-catechin/mg for endosperm and embryo and 16.0 - 16.7  $\mu\text{g}$  ( $\pm$ )-catechin/mg for “pericarb” (pericarp) and testa.<sup>21</sup> *Triticum aestivum* grain may be contaminated with mycotoxins such as trichothecenes and zearaleones from toxin-producing fungi and molds.<sup>14</sup>

#### Triticum Vulgare (Wheat) Bran Extract

Descriptions of the compositions of 4 different *Triticum Vulgare* (Wheat) Bran Extract products are summarized in Table 3.<sup>18</sup>

#### Triticum Vulgare (Wheat) Flour Extract

The yield of non-starch lipids, neutral lipids, glycolipids, and phospholipids of wheat flour extract was dependent on the solvent system used.<sup>22</sup> Total lipids ranged from 6.30 - 9.76 g fatty acid/kg flour, neutral lipids ranged from 3.76 - 5.43 g fatty acid/kg flour, glycolipids ranged from 1.09 - 2.98 g fatty acid/kg flour, and phospholipids ranged from 0.78 - 1.85 g fatty acid/kg flour. The five major fatty acids present in the lipid classes extracted from wheat flour were palmitic acid, stearic acid, oleic acid, linoleic acid, and linolenic acid.

#### Triticum Vulgare (Wheat) Germ

Wheat germ contains about 10% - 15% lipids, 26% - 35% protein, 17% sugar, 1.5% - 4.5% fiber, and about 4% minerals.<sup>16</sup> Bioactive constituents include tocopherols (300 - 740 mg/kg dry matter), phytosterols (24 - 50 mg/kg), policosanols (10 mg/kg), carotenoids (4 - 38 mg/kg), thiamin (15 - 23 mg/kg), and riboflavin (6 - 10 mg/kg).

#### Triticum Vulgare (Wheat) Germ Extract

Analyses of an aqueous mixture containing 3.5% - 12% *Triticum Vulgare* (Wheat) Germ Extract determined that the heavy metals content was below the acceptability threshold of 0.5 ppm, with cobalt, mercury, and lead not quantifiable.<sup>23</sup> No pesticides were detected in this test material.

#### Triticum Vulgare (Wheat) Kernel Flour

There are four classes of protein in wheat flour: globulins, albumins, gliadins, and glutenins.<sup>19</sup> Gliadins and glutenins are components of gluten. Refined wheat flours may contain approximately 13 g/100 g dry weight protein, 0.8 - 0.9 g/100 g dry weight lipids, 85 g/100 g dry weight carbohydrates, and 9.5 g/100 g dry weight gluten.<sup>13</sup> Whole wheat flour contains approximately 15 g/100 g dry weight protein, 1.5 g/100 g dry weight lipids, 82 g/100 g dry weight carbohydrates, and 7 g/100 g dry weight gluten. Wheat flour may be contaminated with foodborne bacteria and fungi, including *Salmonella* spp., *Escherichia coli*, and *Bacillus cereus*. Wheat flour may also be contaminated with mycotoxins such as aflatoxins and ochratoxin A.

#### Triticum Vulgare (Wheat) Protein

Wheat proteins are classified into gluten and non-gluten proteins.<sup>24</sup> The gluten proteins are comprised of gliadins and high and low molecular weight glutenins. The non-gluten proteins are comprised of albumins and globulins.

#### Triticum Vulgare (Wheat) Sprout Extract

A hydroalcoholic extract of wheat sprouts was determined to be composed of hydrophilic compounds, such as peptides, and hydrophobic compounds, such as polyphenols.<sup>25</sup> Further extraction with 90% acetone resulted in the precipitation of 70% glutathione. Phospholipid classes identified from hydroalcoholic extract of wheat sprouts included phosphatidylethanolamine, phosphatidylcholine, phosphatidic acid, phosphatidylinositol, and lysophosphatidylcholine.<sup>26</sup> These phospholipids were characterized by a high content of essential fatty acids ( $\alpha$ -linoleic acid and  $\alpha$ -linolenic acid).

Wheatgrass is reported to contain chlorophyll, flavonoids, several varieties of vitamins including vitamins C and E, choline, minerals, enzymes, indoles, and a number of amino acids.<sup>17</sup>

## Wheat Germ Glycerides

*Wheat Germ Glycerides contain 30% to 40% monoglycerides with di- and triglycerides of mixed fatty acids.*<sup>4</sup>

### USE

#### **Cosmetic**

The safety of the cosmetic ingredients addressed in this assessment is evaluated based on data received from the US Food and Drug Administration (FDA) and the cosmetics industry on the expected use of these ingredients in cosmetics. Use frequencies of individual ingredients in cosmetics are collected from manufacturers and reported by cosmetic product category in the FDA Voluntary Cosmetic Registration Program (VCRP) database. Use concentration data are submitted by the cosmetic industry in response to a survey, conducted by the Personal Care Products Council (Council), of maximum reported use concentrations by product category.

According to 2020 VCRP survey data, *Triticum Aestivum* (Wheat) Germ Extract has the most reported uses in cosmetic products with a total of 293 formulations; the majority of the uses are in leave-on skin care products (Table 4).<sup>27</sup> *Triticum Aestivum* (Wheat) Seed Extract has the second greatest reported number of uses in this safety assessment with 166 formulations; the majority of the uses are also in leave-on skin care products. All other in-use ingredients are reported to be used in significantly fewer formulations. The results of the concentration of use surveys conducted by the Council in 2017 and in 2019 indicate that *Triticum Vulgare* (Wheat) Germ Extract has the highest concentration of use in a leave-on formulation; it is used at up to 13% in face powders.<sup>28,29</sup> The maximum concentrations of use for the remaining ingredients are much lower, with the next highest concentration of use reported for products resulting in leave-on dermal exposure is 0.6% in *Triticum Aestivum* (Wheat) Germ Extract in “other” skin care preparations.

For the three ingredients that were previously reviewed and re-reviewed by the Panel, uses for Wheat Germ Glycerides have decreased considerably according to 2020 VCRP data, with 41 reported uses (37 of which are in leave-on products), down from 128 uses (mostly in lipsticks) reported in 2001 (Table 5).<sup>3,27</sup> The number of uses for the other two ingredients have only increased slightly. The maximum concentration of use range for Wheat Germ Glycerides in 2001 was 0.001% to 25%, with 25% reported in lipsticks; in 2019, the maximum concentration of use range for this ingredient was 0.041% to 0.2%, with 0.2% reported in leave-on face and neck products.<sup>3,29</sup> The ingredients not in use according to the VCRP and industry survey are listed in Table 6.

Wheat-derived ingredients may be used in products that can be incidentally ingested or come into contact with the eyes or mucous membranes; for example, *Triticum Vulgare* (Wheat) Germ Extract is reported to be used in lipstick at up to 0.13%, *Triticum Vulgare* (Wheat) Protein is reported to be used in eye lotion at up to 0.16%, and *Triticum Vulgare* (Wheat) Bran is reported to be used in “other” personal cleanliness products at up to 0.61%.<sup>28</sup>

Additionally, some of the ingredients are used in cosmetic sprays and could possibly be inhaled; for example, *Triticum Vulgare* (Wheat) Germ Extract is reported to be used at up to 0.32% in hair spray and up to 0.11% in spray deodorant.<sup>28</sup> In practice, 95% to 99% of the droplets/particles released from cosmetic sprays have aerodynamic equivalent diameters > 10 µm, with propellant sprays yielding a greater fraction of droplets/particles < 10 µm compared with pump sprays.<sup>30,31</sup> Therefore, most droplets/particles incidentally inhaled from cosmetic sprays would be deposited in the nasopharyngeal and thoracic regions of the respiratory tract and would not be respirable (i.e., they would not enter the lungs) to any appreciable amount.<sup>32,33</sup> There is some evidence indicating that deodorant spray products can release substantially larger fractions of particulates having aerodynamic equivalent diameters in the range considered to be respirable.<sup>32</sup> However, the information is not sufficient to determine whether significantly greater lung exposures result from the use of deodorant sprays, compared to other cosmetic sprays. *Triticum Vulgare* (Wheat) Germ Extract is reportedly used in face powders at concentrations up to 13% and could possibly be inhaled.<sup>28</sup> Conservative estimates of inhalation exposures to respirable particles during the use of loose powder cosmetic products are 400-fold to 1000-fold less than protective regulatory and guidance limits for inert airborne respirable particles in the workplace.<sup>34-36</sup>

The wheat-derived ingredients described in this report are not restricted from use in any way under the rules governing cosmetic products in the European Union.<sup>37</sup>

#### **Non-Cosmetic**

The FDA requires allergen labeling when major allergens, such as wheat, are included in food.<sup>38</sup> A major food allergen is an ingredient from a food or food group, such as wheat, that contains protein derived from the food.

*Triticum vulgare* is reported to be used extensively in traditional medicine to treat decubitus ulcers, burns, scarring delays, dystrophic diseases, and to accelerate tissue repair.<sup>39</sup> It has been investigated for anti-inflammatory properties.

Bran (source not specified) is an over-the-counter (OTC) laxative drug product.<sup>40</sup> Wheat germ is an active ingredient in weight control drug products; however, based on evidence currently available, there are inadequate data to establish general recognition of the safety and effectiveness of this ingredient for this specified use (21CFR§310.545). Wheat gluten is generally recognized as safe (GRAS) in the US for food (21CFR§184.1322).

Wheat leaf extract has been studied for therapeutic benefits for chronic fatigue syndrome, and for its immunological, anti-oxidative, and anti-cancer activities.<sup>17,41</sup> Wheat sprout extract has been studied for its antioxidant content and potential use as a food ingredient and in cancer treatments.<sup>25,26,42</sup> Wheat germ is used as a food supplement and an ingredient in several food products,<sup>16</sup> and wheat germ extract has also been studied for use in cancer prevention and treatment.<sup>43</sup>

### **TOXICOKINETIC STUDIES**

No relevant toxicokinetics studies on wheat-derived ingredients were found in the public literature, and unpublished data were not submitted. In general, toxicokinetics data are not expected to be found on botanical ingredients because each botanical ingredient is a complex mixture of constituents.

### **TOXICOLOGICAL STUDIES**

Most of the wheat-derived ingredients that are reviewed in this safety assessment are found in the foods that are consumed daily, and daily exposure from food use would result in much larger systemic exposures than those from use in cosmetic products. The potential for systemic exposure from the absorption of these ingredient through the skin is much less than the potential for systemic exposure from absorption through oral exposures. This is because the rates of absorption and metabolism of these ingredients in the skin are expected to be negligible compared to the corresponding rates in the digestive tract. Thus, the potential for systemic effects, other than sensitization, is not discussed in detail in this report.

### **DEVELOPMENTAL AND REPRODUCTIVE TOXICITY (DART) STUDIES**

#### **Oral**

##### ***Triticum Vulgare (Wheat) Sprout Extract***

The effects of the water extract of wheat sprouts on spermatozoa was investigated using male BALB/c mice.<sup>44</sup> The mice were treated with benzo[a]pyrene (B[a]P) in corn oil (100 mg/kg bw) for 5 consecutive days via intraperitoneal injections. A control group of 19 mice only received corn oil (0.5 ml) and two groups of 3 mice only received wheat sprout extract (40 and 70 ml). Nineteen animals received the wheat sprout extract in parallel with the B[a]P treatment (3 mice received 40 ml extract, 10 mice received 70 ml extract, and 6 mice received 140 ml extract, in 18 equal doses by gavage) starting 3 days before the B[a]P treatment. Twenty-eight mice received just B[a]P. The sperm of the treated mice were examined 5 weeks after the B[a]P treatment. The mice treated only with B[a]P had an incidence of 61.1% for abnormally-shaped sperm heads. The corn oil control group only had sperm abnormalities observed in 1.93%. The wheat sprout extract alone did not enhance the level of sperm abnormalities in comparison with the corn oil controls. The simultaneous treatment of B[a]P and wheat sprout extract resulted in a decrease in the percentage of abnormally-shaped sperm heads when compared to the group that received just B[a]P, but not in a dose-dependent manner.

### **GENOTOXICITY**

#### **In Vitro**

##### ***Triticum Vulgare (Wheat) Bran Extract***

The genotoxicity potential of a wheat bran extract that was highly-enriched in arabinoxylan-oligosaccharides was assessed in an Ames test using *S. typhimurium* strains TA 98, TA 100, TA 1535, and TA 1537 and *E. coli* strain WP2uvrA, with and without metabolic activation.<sup>45</sup> The extract was dissolved at 100 mg/ml in distilled water and was tested at 5, 15.8, 50, 158, 500, 1581, and 5000 µg/plate. Strain-specific positive and negative control chemicals were used and yielded expected results. Wheat bran extract was not mutagenic at up to 5000 µg/plate.

The same researchers assessed the clastogenic potential of the same wheat bran extract in a chromosome aberration assay using Chinese hamster lung fibroblast V79 cells.<sup>45</sup> The cells were treated for 3 h or 20 h with the extract at up to 5000 µg/ml, with and without metabolic activation. No biologically significant increases in the number of cells showing structural chromosome aberrations were observed. The wheat bran extract was not clastogenic in this assay.

##### ***Wheat Germ Glycerides***

The genotoxicity potential of Wheat Germ Glycerides in ethanol was assessed in an Ames test using *S. typhimurium* strains TA 98, TA 100, TA 102, TA 1535, and TA 1537, with and without metabolic activation.<sup>46</sup> Strain-specific positive and negative control chemicals were used and yielded expected results. Wheat Germ Glycerides was not mutagenic when tested at up to 5000 µg/plate.

## ANTI-MUTAGENICITY

### **In Vitro**

#### *Triticum Vulgare (Wheat) Sprout Extract*

The ability of the S-30 fraction (decanted, incubated supernatant fluid) of the water extract of wheat sprouts to inhibit mutagenicity was assessed in an Ames test using *S. typhimurium* strain TA 98, with metabolic activation and in the presence of 7,12-dimethyl benz[a]anthracene (DMBA; 50 µg/plate).<sup>44</sup> The S-30 fraction was prepared from thawed plants by squeezing out the juice and centrifuging it for 30 min at 30,000 g before removing proteins with heat, concentrating by vacuum, and sterilizing. A decrease in the number of revertants was observed. No further details were provided.

### **In Vivo**

#### *Triticum Vulgare (Wheat) Sprout Extract*

The same research group from the above study investigated the effects of the S-30 fraction of the wheat sprout extract on the ability of BALB/c mouse skin enzymes to activate DMBA to mutagenic metabolites.<sup>44</sup> The S-9 fraction (added to in vitro systems to simulate metabolic capability) from the mouse skin that received the wheat extract subcutaneously for 4 days (0.1 ml/day) still activated DMBA (single painting; 100 µg/mouse) to mutagenic metabolites for the *S. typhimurium* strain TA 98, but the ability was 20% lower than that of the S-9 fraction from mice that did not receive the wheat sprout extract. No further details were provided.

## CARCINOGENICITY

### **Co-Carcinogenicity**

#### *Triticum Vulgare (Wheat) Bran*

In a 31-week dietary study, a group of 12 male Sprague-Dawley rats received 20% wheat bran as a dietary supplement during and after administration of 1,2-dimethylhydrazine (a carcinogen) via 13 weekly subcutaneous injections.<sup>47</sup> Another group of 12 rats was fed the wheat bran in conjunction with administration of the carcinogen, and then switched to the control diet after 13 weeks. A third group of 11 rats received a fiber-free (control) diet during the administration of the carcinogen, and then switched to the wheat bran diet after 13 weeks. The control group (12 rats) received a fiber-free diet throughout the 31 weeks. In both groups that received wheat bran in conjunction with administration of the carcinogen, small intestinal or colonic tumor yield was significantly greater, with benign and malignant tumors incidences increased by 3.4-fold ( $p < 0.005$ ), adenoma incidences increased by 3.5-fold ( $p < 0.025$ ), and adenocarcinomas incidences increased by 3.25-fold ( $p < 0.05$ ) when compared to rats that received a control diet. Rats that received wheat bran only after carcinogen exposure had a reduced number of benign adenomas (71.4%;  $p < 0.025$ ) when compared to the rats that received control diet. The rats fed the wheat bran during carcinogen administration and then switched to the control diet had the highest tumor yield, with 4.5 times as many benign and malignant tumors as the control rats ( $p < 0.05$ ), and at least 6 times as many adenomas as any of the other dietary groups ( $p < 0.05$ ).

### **Tumor/Anti-Tumor Promotion**

#### *Triticum Aestivum (Wheat) Leaf Extract*

The effects of wheatgrass leaf extract on skin papillomagenesis induced by DMBA and croton oil were investigated in male Swiss albino mice.<sup>48</sup> The mice were divided into five groups of 10: Group 1 was the control group that received a single dose of DMBA (100 µl/50 µl acetone) on shaved dorsal skin followed two weeks later by croton oil (1% in 100 µl acetone) three times a week for a total of 16 weeks; Group 2 (pre-group) received wheatgrass leaf extract orally (20 ml/kg bw) for 7 days prior to the application of DMBA and croton oil in the manner that Group 1 received these; Group 3 (peri-group) was treated with DMBA and then received wheatgrass leaf extract for 15 days prior to receiving the croton oil, as in Group 1; Group 4 (post-group) was treated with DMBA as in Group 1 and received wheatgrass leaf extract at the same time the croton oil treatment started; and Group 5 received the wheatgrass leaf extract seven days prior to and after the treatment with DMBA and through the treatment of croton oil until experiment end (total number of exposure days not stated). Tumor incidence, yield, and burden in Groups 2-5 were significantly decreased as compared to Group 1. The average latent period was increased in Groups 2-5, with the longest latent period being observed in Group 5.

#### *Triticum Vulgare (Wheat) Sprout Extract*

The tumorigenic effects of wheat sprout extract (S-30 fraction) were investigated using 8-week-old male BALB/c mice.<sup>44</sup> Skin papillomas were initiated by painting the skin of mice with an acetone solution of DMBA (100 µg/mouse). Twice weekly treatments of croton oil dissolved in acetone (10 µl) was used for papilloma growth promotion. The mice received the promoter for 22 weeks. One group of mice (n = 38) received 10 successive subcutaneous injections of wheat sprout extract (0.1 ml) starting on day 3 before DMBA treatment) with the croton oil promoter, while another group (n = 8) received the extract during the promotion period without the croton oil promoter. Further groupings involved giving mice the wheat sprout extract (0.1 ml) subcutaneously twice a week for 22 weeks without the croton oil promoter during both the initiator and promoter phases (n =

13). Wheat sprout extract, when injected subcutaneously for 10 days during carcinogenesis initiation in mice, shortened the latency period from 9 to 4 weeks and increased the number of skin papillomas by 4-fold. When the extract was applied to mice treated with DMBA, but did not have croton oil promotion, only one mouse developed papillomas during the 24 weeks. Controls where the extract was injected alone without initiation or promotion did not produce skin papillomas (n = 17). The authors concluded that wheat sprout extract did not have initiating or promoting properties.

## **IRRITATION AND SENSITIZATION**

### **Dermal Irritation**

#### **Animal**

##### **Triticum Vulgare (Wheat) Germ Extract**

In a cutaneous tolerance study of an aqueous solution containing 12% Triticum Vulgare (Wheat) Germ Extract, three rabbits received the test material as a single dose (0.5 ml) on the skin in accordance with Organization for Economic Co-operation and Development (OECD) test guideline (TG) 404.<sup>23</sup> A slight to clear erythema that was reversible within 72 h was observed in all animals. It was concluded that the test material was not an irritant to the skin. No additional details were provided.

##### **Wheat Germ Glycerides**

*Mild irritation was observed on abraded rabbit skin following dermal application of different lots of Wheat Germ Glycerides.<sup>4</sup> Minimal skin irritation was observed in rabbits that received dermal applications of Wheat Germ Glycerides in several different cosmetic formulations. Concentrations were not provided for any of these studies.*

### **Dermal Sensitization**

#### **Animal**

##### **Wheat Germ Glycerides**

*Wheat Germ Glycerides (0.1% solution in olive oil) was evaluated for sensitization in a guinea pig sensitization study.<sup>4</sup> The treatment group comprised 6 guinea pigs that received an initial 0.05 ml intracutaneous injection and 9 subsequent 0.1 ml injections during the induction phase followed by a 2-week rest period. The animals then received a 0.05 ml challenge injection. The test material was not a skin sensitizer.*

#### **Human**

##### **Triticum Vulgare (Wheat) Germ Extract**

A human repeated insult patch test (HRIPT) of a face and body powder containing 13% Triticum Vulgare (Wheat) Germ Extract was conducted in 105 subjects.<sup>49</sup> The test material (200 mg) was applied to the test sites as supplied. The patches were partially occlusive and 2 cm<sup>2</sup> in area. Nine sequential 24-h induction applications and 2 concurrent 24-h challenge applications were made, with the challenge patches on the initial induction site and a naïve site. Challenge sites were read at 24 and 48 h post-application. No irritation or sensitization was observed.

##### **Triticum Vulgare (Wheat) Gluten and Wheat Germ Glycerides**

*Mascara-base products containing up to 1% Triticum Vulgare (Wheat) Gluten were not dermal irritants or sensitizers in HRIPTs of up to 202 subjects or in in-use studies (number of subjects not reported).<sup>4</sup> A lipstick base containing up to 2% Wheat Germ Glycerides produced reactions in 8 out of 1154 subjects: reactions were considered to be irritation. Wheat Germ Glycerides were not dermal irritants or sensitizers in HRIPTs of lipstick products and blushers at concentrations up to 0.5%. No adverse reactions were reported in in-use studies of a lipstick base containing 2% Wheat Germ Glycerides in 149 subjects.*

## **OCULAR IRRITATION STUDIES**

##### **Triticum Vulgare (Wheat) Gluten and Wheat Germ Glycerides**

*Mild circumcorneal injection and mild discharge were noted after testing a mascara base containing 1% Triticum Vulgare (Wheat) Gluten in rabbit eyes: these changes were thought to be the results of foreign objects in the eye and drying of the formulation.<sup>4</sup> Complete recovery occurred within 72 h. Minimal to no ocular irritation was observed in rabbit studies with undiluted Wheat Germ Glycerides and in formulations containing up to 2% Wheat Germ Glyceride.*

##### **Triticum Vulgare (Wheat) Germ Extract**

In an ocular tolerance study of an aqueous solution containing 12% Triticum Vulgare (Wheat) Germ Extract, three rabbits received the test material as a single dose (0.1 ml) instilled into the eye in accordance with OECD TG 405.<sup>23</sup> Mild irritation was observed that was reversible within 72 h. It was concluded that the test material was not an irritant to the rabbit eye. No additional details were provided.



## OCCUPATIONAL EXPOSURES

Work-related sensitization (IgE-mediated) to wheat flour and grain dusts has been reported in bakery workers.<sup>50-54</sup> Commonly known as baker's asthma, reactions are often preceded by rhinitis and other respiratory symptoms, with concomitant skin symptoms such as contact urticaria and hand eczema. Atopy and sensitization to flour and/or enzyme (e.g.  $\alpha$ -amylase of fungal origin) occurs frequently.<sup>53,54</sup> Aside from wheat, baker's asthma may also be caused by molds, yeast, eggs, sesame seeds, nuts, and insects. Skin-prick testing, skin biopsies, and radioallergosorbent tests (RAST) have been utilized to identify and analyze the reactions observed in bakery workers.<sup>50-54</sup>

## EPIDEMIOLOGY OF IMMUNE-MEDIATED GLUTEN AND WHEAT REACTIONS

Celiac disease affects approximately 1% of the population worldwide, including the US, with variations between countries.<sup>55-57</sup> Wheat allergy affects between 0.5% to 9% of the population globally, with about 0.4% of adults in the US diagnosed with the allergy.<sup>56,58,59</sup> Children have a higher prevalence to food allergy to wheat compared to adults, and respiratory IgE mediated allergies to wheat (i.e. baker's asthma or rhinitis) are often associated as occupational disease with the incidence in bakers ranging at 5% - 10% for asthma and 15% - 20% for rhinitis.<sup>60</sup> Bronchial reactivity to inhaled wheat proteins in adults with food allergy is very rare but has been documented.<sup>61</sup>

## SUMMARY

Most of the 27 wheat-derived ingredients detailed in this safety assessment are reported to function in cosmetics as skin conditioning agents, while some are reported to have other functions, such as abrasives, absorbents, antioxidants, bulking agents, film formers, flavoring agent, hair conditioning agents and viscosity increasing agents. Functions such as skin bleaching agent (for *Triticum Vulgare* (Wheat) Germ Extract) are not considered cosmetic functions in the US and, therefore, are not addressed in this assessment.

Yields of constituents in wheat have been found to be dependent on extraction methods and growing conditions such as soil composition, climate, duration of growth period, and height at harvest. Additionally, different plant parts have different constituent compositions. Wheat grain may be contaminated by mycotoxins, such as aflatoxins, trichothecenes, and zearalenones. Wheat flour may be contaminated with mycotoxins and foodborne bacteria and fungi, such as *Salmonella* spp., *E. coli*, and *B. cereus*.

According to 2020 VCRP survey data, *Triticum Aestivum* (Wheat) Germ Extract has the most reported uses in cosmetic products with a total of 293 formulations; the majority of the uses are in leave-on skin care products. *Triticum Aestivum* (Wheat) Seed Extract has the second greatest reported number of uses in this safety assessment with 166 formulations; the majority of the uses are also in leave-on skin care products. All other in-use ingredients are reported to be used in significantly fewer formulations. The results of the concentration of use survey conducted by the Council indicate that *Triticum Vulgare* (Wheat) Germ Extract has the highest maximum concentration of use in a leave-on formulation; it is used at up to 13% in face powders. The concentrations of use for the remaining ingredients are much lower; the next highest maximum concentration of use reported for products resulting in leave-on dermal exposure is 0.6% in *Triticum Aestivum* (Wheat) Germ Extract in "other" skin care preparations.

Wheat is considered a major food allergen and is required by the FDA to be labeled as such when included in food. *Triticum vulgare* is reported to be used extensively in traditional medicine and it has been investigated for anti-inflammatory properties. Wheat bran is an OTC laxative drug product. Wheat leaf extract and wheat germ extract have been studied for various therapeutic effects and for use in food supplements.

Most of the wheat-derived ingredients that are reviewed in this safety assessment are found in foods consumed daily the world over. The potential for systemic exposure from the absorption of these ingredient through the skin is much less than the potential for systemic exposure from absorption through oral exposures. This is because the rates of absorption and metabolism of these ingredients in the skin are expected to be negligible compared to the corresponding rates in the digestive tract; and, the systemically available dose of these ingredients, even with theoretically complete absorption from cosmetic use, would be very small compared to that available from consumption.

Wheat sprout extract (40 and 70 ml) alone did not enhance the level of sperm abnormalities in mice in comparison with controls. Simultaneous treatment of B[a]P and wheat sprout extract resulted in a decrease in the percentage of abnormally-shaped sperm heads when compared to the group that received just B[a]P, but not in a dose-dependent manner.

Wheat bran extract was not mutagenic in an Ames test at up to 5000  $\mu\text{g}/\text{plate}$ , nor was it clastogenic in Chinese hamster lung fibroblasts in a chromosome aberration assay at up to 5000  $\mu\text{g}/\text{ml}$ . In anti-mutagenicity studies, a decrease in the number of revertants was observed following exposure to the S-30 fraction of a water extract of wheat sprouts in an Ames test with metabolic activation and DMBA. A decrease in mutagenic metabolites was observed in BALB/c mouse skin following subcutaneous exposure to the same wheat sprout extract (0.1 ml/day) and DMBA.

Tumor incidences were increased in a 31-week study of rats that received a dietary supplement containing 20% wheat bran during administration of a carcinogen for 13 weeks when compared to rats that received a control diet. Rats that received the

wheat bran after carcinogen exposure had a reduced number for benign adenomas when compared to controls. Wheat sprout extract (0.1 ml) was not an initiator or a promoter in mice that received the extract subcutaneously, were induced with DMBA, and/or promoted with croton oil in a 24-week study. Anti-tumorigenic effects have been reported in an oral study with wheatgrass extract (20 ml/kg bw) in mice that were induced with DMBA and promoted with croton oil.

No irritation or sensitization was observed during an HRIPT of a face and body powder containing 13% *Triticum Vulgare* (Wheat) Germ Extract that was conducted in 105 subjects. No dermal or ocular irritation to 12% *Triticum Vulgare* (Wheat) Germ Extract was observed in rabbits in a cutaneous tolerance study and an ocular tolerance study, respectively.

Work-related sensitization has been reported in bakery workers. Symptoms include rhinitis, asthma, contact urticaria, and hand eczema. Celiac disease and wheat allergy affect populations worldwide. Children have a higher prevalence to food allergy to wheat compared to adults, and respiratory IgE mediated allergies to wheat (i.e. baker's asthma or rhinitis) are often associated as occupational disease. Bronchial reactivity to inhaled wheat proteins in adults with food allergy is very rare but has been documented.

No relevant toxicokinetic studies were found in the published literature; however, in general, toxicokinetics data are not expected to be found on botanical ingredients because each botanical ingredient is a complex mixture of constituents.

## DISCUSSION

The Panel reviewed the botanical ingredients derived from the wheat plants *Triticum aestivum*, *Triticum monococcum*, *Triticum spelta*, *Triticum turgidum durum*, and *Triticum vulgare*. The accepted scientific name for both *Triticum vulgare* and *Triticum spelta* is *Triticum aestivum*. The Panel expressed concern about pesticide residues, heavy metals, and other plant species that may be present in botanical ingredients, and stressed that the cosmetics industry should continue to use current good manufacturing practices (cGMPs) to limit impurities.

While aflatoxin has been detected in wheat grain and flour, the Panel believes that aflatoxin should not be present in wheat-derived ingredients that are derived from *Triticum aestivum*, *Triticum monococcum*, *Triticum spelta*, *Triticum turgidum durum*, or *Triticum vulgare*. The Panel has adopted the United States Department of Agriculture (USDA) designation of  $\leq 15$  ppb as corresponding to "negative" aflatoxin content.

The Panel noted that it had previously concluded that Hydrolyzed Wheat Protein and Hydrolyzed Wheat Gluten were safe for use in cosmetics when formulated to restrict peptides to an average molecular weight of 3500 Daltons (Da) or less. This conclusion was in response to reports of type 1 (IgE-mediated) immediate hypersensitivity reactions that occurred in sensitized individuals following exposure to cosmetic products that contained one of these two ingredients with molecular weights greater than this limit. However, based on the available information, none of the wheat-derived ingredients in this report are hydrolyzed and most are not even proteins. Coupled with lack of reports to the contrary or experience with such reactions to these ingredients in the clinical setting, concern of over such reactions to these ingredients was mitigated. If the protein ingredients in this report are hydrolyzed in processing, then the Panel needs to be made aware of these methods of manufacturing to further assess the safety of these ingredients.

Some wheat-derived ingredients were reported to be used in spray and powder products that could possibly be inhaled. For example, *Triticum Vulgare* (Wheat) Germ Extract is reported to be used at up to 0.32% in hair spray, at up to 0.11% in spray deodorant, and at up to 13% in face powders. The Panel noted that in aerosol products, 95% – 99% of droplets/particles would not be respirable to any appreciable amount. Furthermore, droplets/particles deposited in the nasopharyngeal or bronchial regions of the respiratory tract present no toxicological concerns for these ingredients. Coupled with the small actual exposure in the breathing zone and the concentrations at which the ingredients are used, the available information indicates that incidental inhalation would not be a significant route of exposure that might lead to local respiratory or systemic effects. A detailed discussion and summary of the Panel's approach to evaluating incidental inhalation exposures to ingredients in cosmetic products is available at <https://www.cir-safety.org/cir-findings>.

At its June 2020 meeting, the Panel determined that data are insufficient to determine safety. The additional data needed to determine the safety of the wheat-derived ingredients as used in cosmetics are:

- Method of manufacturing data
- Dermal irritation and sensitization data at a test concentration of 13% for *Triticum Vulgare* (Wheat) Sprout Extract

## CONCLUSION

The Expert Panel for Cosmetic Ingredient Safety concluded that the available data are insufficient to make a determination the following 27 ingredients under the intended conditions of use in cosmetic formulations:

Triticum Aestivum (Wheat) Flour Lipids	Triticum Vulgare (Wheat) Flour Lipids
Triticum Aestivum (Wheat) Germ Extract	Triticum Vulgare (Wheat) Germ
Triticum Aestivum (Wheat) Leaf Extract*	Triticum Vulgare (Wheat) Germ Extract
Triticum Aestivum (Wheat) Peptide*	Triticum Vulgare (Wheat) Germ Powder*
Triticum Aestivum (Wheat) Seed Extract	Triticum Vulgare (Wheat) Germ Protein
Triticum Monococcum (Wheat) Seed Extract	Triticum Vulgare (Wheat) Gluten
Triticum Monococcum (Wheat) Stem Water*	Triticum Vulgare (Wheat) Gluten Extract
Triticum Spelta Seed Water*	Triticum Vulgare (Wheat) Kernel Flour
Triticum Turgidum Durum (Wheat) Seed Extract*	Triticum Vulgare (Wheat) Protein
Triticum Vulgare/Aestivum (Wheat) Grain Extract*	Triticum Vulgare (Wheat) Seed Extract
Triticum Vulgare (Wheat) Bran	Triticum Vulgare (Wheat) Sprout Extract
Triticum Vulgare (Wheat) Bran Extract	Triticum Vulgare (Wheat) Straw Water*
Triticum Vulgare (Wheat) Bran Lipids	Wheat Germ Glycerides
Triticum Vulgare (Wheat) Flour Extract	

*\*There are currently no uses reported for these ingredients.*

## TABLES

**Table 1. Definitions and functions of the ingredients in this safety assessment.<sup>1</sup>**

<b>Ingredient/CAS No.</b>	<b>Definition</b>	<b>Function</b>
Triticum Aestivum (Wheat) Flour Lipids	Triticum Aestivum (Wheat) Flour Lipids is a mixture of lipids derived from the flour of <i>Triticum aestivum</i> .	Skin-conditioning agent – misc.
Triticum Aestivum (Wheat) Germ Extract	Triticum Aestivum (Wheat) Germ Extract is the extract of the germs of <i>Triticum aestivum</i> .	Antioxidant; skin-conditioning agent – misc.
Triticum Aestivum (Wheat) Leaf Extract	Triticum Aestivum (Wheat) Leaf Extract is the extract of the leaves of the wheat, <i>Triticum aestivum</i> .	Antioxidant
Triticum Aestivum (Wheat) Peptide	Triticum Aestivum (Wheat) Peptide is the di-/tri-peptide fraction isolated from the protein of <i>Triticum aestivum</i> by ultra-membrane filtration.	Film former; hair conditioning agent; skin-conditioning agent – misc.
Triticum Aestivum (Wheat) Seed Extract	Triticum Aestivum (Wheat) Seed Extract is the extract of the seeds of the wheat, <i>Triticum aestivum</i> .	Hair conditioning agent; skin-conditioning agent – misc.
Triticum Monococcum (Wheat) Seed Extract	Triticum Monococcum (Wheat) Seed Extract is the extract of the seeds of the wheat, <i>Triticum monococcum</i> . [ <i>Triticum monococcum</i> is also known as eikorn and is native to eastern Europe and western Asia]. <sup>12</sup>	Skin-conditioning agent – misc.
Triticum Monococcum (Wheat) Stem Water	Triticum Monococcum (Wheat) Stem Water is the aqueous solution of the steam distillates obtained from the stems of <i>Triticum monococcum</i> .	Flavoring agent
Triticum Spelta Seed Water	Triticum Spelta Seed Water is the aqueous solution of the steam distillates obtained from the seeds of <i>Triticum spelta</i> . *	Skin-conditioning agent – misc.
Triticum Turgidum Durum (Wheat) Seed Extract	Triticum Turgidum Durum (Wheat) Seed Extract is the extract of the seeds of <i>Triticum turgidum durum</i> .	Skin-conditioning agent – misc.
Triticum Vulgare/Aestivum (Wheat) Grain Extract	Triticum Vulgare/Aestivum (Wheat) Grain Extract is the extract of the grains of <i>Triticum vulgare</i> and <i>Triticum aestivum</i> .	Hair conditioning agent; skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Bran	Triticum Vulgare (Wheat) Bran is the broken coat material of grains of wheat, <i>Triticum vulgare</i> .*	Abrasive; bulking agent
Triticum Vulgare (Wheat) Bran Extract 84012-44-2	Triticum Vulgare (Wheat) Bran Extract is the extract of the bran of <i>Triticum vulgare</i> .*	Skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Bran Lipids	Triticum Vulgare (Wheat) Bran Lipids is the cyclohexane extract of Triticum Vulgare (Wheat) Bran.*	Skin-conditioning agent - occlusive
Triticum Vulgare (Wheat) Flour Extract	Triticum Vulgare (Wheat) Flour Extract is the extract of the powder obtained by grinding wheat, <i>Triticum vulgare</i> .*	Skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Flour Lipids	Triticum Vulgare (Wheat) Flour Lipids is a mixture of lipids derived from the flour of <i>Triticum vulgare</i> .*	Skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Germ	Triticum Vulgare (Wheat) Germ is the natural product obtained from the embryo of the wheat kernel separated in milling.*	Skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Germ Extract 84012-44-2	Triticum Vulgare (Wheat) Germ Extract is the extract of the germ of <i>Triticum vulgare</i> .*	Skin bleaching agent; skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Germ Powder	Triticum Vulgare (Wheat) Germ Powder is the powder obtained from the dried, ground wheat germ, <i>Triticum vulgare</i> .*	Abrasive; absorbent; bulking agent; viscosity increasing agent - aqueous
Triticum Vulgare (Wheat) Germ Protein	Triticum Vulgare (Wheat) Germ Protein is a protein obtained from the germ of the wheat, <i>Triticum vulgare</i> .*	Hair conditioning agent; skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Gluten 8002-80-0	Triticum Vulgare (Wheat) Gluten is a protein substance which is intermixed with the starch portion in the endosperm of the wheat, <i>Triticum vulgare</i> .*	Binder; hair conditioning agent; skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Gluten Extract	Triticum Vulgare (Wheat) Gluten Extract is the extract of Triticum Vulgare (Wheat) Gluten.*	Skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Kernel Flour	Triticum Vulgare (Wheat) Kernel Flour is the milled flour obtained from the finely ground kernels of wheat, <i>Triticum vulgare</i> .*	Abrasive; bulking agent; viscosity increasing agent - aqueous
Triticum Vulgare (Wheat) Protein	Triticum Vulgare (Wheat) Protein is a protein obtained from wheat, <i>Triticum vulgare</i> .*	Film former; hair conditioning agent; skin-conditioning agent – misc.

**Table 1. Definitions and functions of the ingredients in this safety assessment.<sup>1</sup>**

<b>Ingredient/CAS No.</b>	<b>Definition</b>	<b>Function</b>
Triticum Vulgare (Wheat) Seed Extract 84012-44-2	Triticum Vulgare (Wheat) Seed Extract is the extract of the seeds of <i>Triticum vulgare</i> .*	Skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Sprout Extract	Triticum Vulgare (Wheat) Sprout Extract is the extract of the young shoots of <i>Triticum vulgare</i> .*	Skin-conditioning agent – misc.
Triticum Vulgare (Wheat) Straw Water	Triticum Vulgare (Wheat) Straw Water is the aqueous solution of the steam distillates obtained from <i>Triticum vulgare</i> (wheat) straw.*	Skin-conditioning agent – misc.
Wheat Germ Glycerides 68990-07-8	Wheat Germ Glycerides is a mixture of mono-, di-, and triglycerides produced by the transesterification of Triticum Vulgare (Wheat) Germ Oil.	Skin-conditioning agent - emollient

\* The accepted scientific name for both *Triticum vulgare* and *Triticum spelta* is *Triticum aestivum*.<sup>1</sup>

**Table 2. Generic plant part definitions as they apply to wheat-derived ingredients.<sup>1</sup>**

<b>Plant Part</b>	<b>Definition</b>
Bran	The outer hard layers of the grain formed by the fused fruit and seed wall in grains and cereals.
Endosperm	Energy storage tissue inside seeds.
Germ	The embryo in a seed; the part of a seed that can develop into new plant.
Grain	Dry one-seeded fruits produced by grasses, e.g. cereals such as wheat.
Kernel	The grain of a grass.
Leaf	Flattened photosynthetic organs, attached to stems.
Pericarp	Fruit wall.
Seed	A propagating sexual structure resulting from the fertilization of an ovule, formed by embryo, endosperm, or seed coat.
Seed coat	Seed wall; testa; protective outer layer of seed, formed from the outer layers of the ovule
Sprout	Seedling; germinating seed; any new growth of a plant from a stem such as a new branch or a bud
Stem	A slender or elongated structure that supports a plant or a plant part or plant organ.
Straw	The stem of a grass or related families

**Table 3. Composition of 4 different Triticum Vulgare (Wheat) Bran Extract products<sup>18</sup>**

<b>Extract</b>	<b>Glycerin Extract</b>	<b>Propylene Glycol Extract</b>	<b>Propylene Glycol Extract</b>	<b>Soybean Oil Extract</b>
<b>Composition of Mixture</b>	75%-100% glycerin 10%-25% water 5%-10% Triticum Vulgare (Wheat) Bran Extract	75%-100% propylene glycol 10%-25% Triticum Vulgare (Wheat) Bran Extract	95%-100% propylene glycol 1%-5% Triticum Vulgare (Wheat) Bran Extract	75%-100% soybean oil 10%-25% Triticum Vulgare (Wheat) Bran Extract
<b>Solvent of Extraction</b>	glycerin (vegetable origin)	1,2-propylene glycol	1,2-propylene glycol	soybean oil
<b>Preservatives</b>	0.35% potassium sorbate 0.35% sodium benzoate	0.35% potassium sorbate 0.35% sodium benzoate	0.35% potassium sorbate 0.35% sodium benzoate	none
<b>Incidental Ingredients</b>	0.1%-1% lactic acid	none	0.1%-1% lactic acid	none
<b>Bacteriological Controls</b>	100 germs/ml max.	100 germs/ml max.	100 germs/ml max.	100 germs/ml max.
<b>Heavy Metals (solvent)</b>	≤ 5 ppm total	not tested	not tested	not tested



**Table 4. Frequency (2020) and concentration of use (2017, 2019) according to duration and type of exposure for *Triticum aestivum* (Wheat)-derived ingredients.<sup>27-29</sup>**

	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)	# of Uses	Max Conc of Use (%)
	Triticum Vulgare (Wheat) Flour Lipids <sup>e</sup>		Triticum Vulgare (Wheat) Germ <sup>c</sup>		Triticum Vulgare (Wheat) Germ Extract <sup>*</sup>		Triticum Vulgare (Wheat) Germ Protein <sup>c</sup>	
<b>Totals<sup>†</sup></b>	<b>NR</b>	<b>0.00065-0.1</b>	<b>5</b>	<b>NR</b>	<b>40</b>	<b>0.00001-13</b>	<b>60</b>	<b>0.0015-0.03</b>
<b>Duration of Use</b>								
Leave-On	NR	0.00065-0.1	4	NR	35	0.00001-13	12	0.0015-0.03
Rinse Off	NR	NR	1	NR	5	0.001-0.32	48	0.0075
Diluted for (Bath) Use	NR	NR	NR	NR	NR	NR	NR	NR
<b>Exposure Type</b>								
Eye Area	NR	0.00065	NR	NR	2	0.004-0.075	1	0.0075-0.01
Incidental Ingestion	NR	NR	NR	NR	NR	0.13	NR	NR
Incidental Inhalation-Spray	NR	0.05 <sup>a</sup>	3 <sup>b</sup>	NR	19 <sup>a</sup> ; 9 <sup>b</sup>	0.005-0.32; 0.0012-0.025 <sup>a</sup> ; 0.02 <sup>b</sup>	2 <sup>a</sup> ; 7 <sup>b</sup>	NR
Incidental Inhalation-Powder	NR	0.1 <sup>c</sup>	3 <sup>b</sup>	NR	9 <sup>b</sup>	13; 0.02 <sup>b</sup> ; 0.0001-0.2 <sup>c</sup>	7 <sup>b</sup>	0.015; 0.0075-0.03 <sup>c</sup>
Dermal Contact	NR	0.05-0.1	5	NR	37	0.00001-13	13	0.0075-0.03
Deodorant (underarm)	NR	NR	NR	NR	NR	0.1-0.11 <sup>d</sup>	NR	NR
Hair - Non-Coloring	NR	NR	NR	NR	2	0.001-0.32	NR	NR
Hair-Coloring	NR	NR	NR	NR	NR	NR	46	NR
Nail	NR	NR	NR	NR	NR	NR	1	0.0015
Mucous Membrane	NR	NR	NR	NR	NR	0.1-0.13	1	NR
Baby Products	NR	NR	NR	NR	NR	NR	NR	NR
<b>Triticum Vulgare (Wheat) Gluten Extract</b>								
<b>Totals<sup>†</sup></b>	<b>1</b>	<b>NR</b>	<b>93</b>	<b>0.01-0.16</b>	<b>NR</b>	<b>0.34</b>	<b>13</b>	<b>NR</b>
<b>Duration of Use</b>								
Leave-On	1	NR	76	0.01-0.16	NR	0.34	11	NR
Rinse Off	NR	NR	17	NR	NR	NR	2	NR
Diluted for (Bath) Use	NR	NR	NR	NR	NR	NR	NR	NR
<b>Exposure Type</b>								
Eye Area	NR	NR	16	0.16	NR	NR	1	NR
Incidental Ingestion	NR	NR	NR	NR	NR	NR	NR	NR
Incidental Inhalation-Spray	NR	NR	1; 7 <sup>a</sup> ; 34 <sup>b</sup>	NR	NR	NR	2 <sup>a</sup> ; 5 <sup>b</sup>	NR
Incidental Inhalation-Powder	NR	NR	34 <sup>b</sup>	NR	NR	0.34 <sup>c</sup>	5 <sup>b</sup> ; 2 <sup>c</sup>	NR
Dermal Contact	NR	NR	76	0.16	NR	0.34	11	NR
Deodorant (underarm)	NR	NR	NR	NR	NR	NR	NR	NR
Hair - Non-Coloring	1	NR	17	NR	NR	NR	1	NR
Hair-Coloring	NR	NR	NR	NR	NR	NR	1	NR
Nail	NR	NR	NR	0.01	NR	NR	NR	NR
Mucous Membrane	NR	NR	1	NR	NR	NR	NR	NR
Baby Products	NR	NR	NR	NR	NR	NR	3	NR

NR = Not reported; NS = Not yet surveyed

<sup>†</sup> Because each ingredient may be used in cosmetics with multiple exposure types, the sum of all exposure types may not equal the sum of total uses.

<sup>\*</sup> VCRP data was listed generically as Wheat Germ Extract

<sup>a</sup> It is possible these products may be sprays, but it is not specified whether the reported uses are sprays.

<sup>b</sup> Not specified whether a powder or a spray, so this information is captured for both categories of incidental inhalation.

<sup>c</sup> It is possible these products may be powders, but it is not specified whether the reported uses are powders.

<sup>d</sup> 0.11% in a spray deodorant

<sup>e</sup> Listed with the nomenclature Triticum Aestivum in the VCRP database.

**Table 5. Current and historical frequency and concentration according to duration and type of exposure for previously reviewed wheat-derived ingredients.**<sup>3,27,29</sup>

	Triticum Vulgare (Wheat) Gluten*				Triticum Vulgare (Wheat) Kernel Flour*				Wheat Germ Glycerides			
	# of Uses		Max Conc of Use (%)		# of Uses		Max Conc of Use (%)		# of Uses		Max Conc of Use (%)	
	2020	2001	2019	2001	2020	2001	2019	2001	2020	2001	2019	2001
<b>Totals*</b>	<b>21</b>	<b>5</b>	<b>0.0001-0.05</b>	<b>NR</b>	<b>4</b>	<b>NR</b>	<b>NR</b>	<b>NR</b>	<b>41</b>	<b>128</b>	<b>0.041-0.2</b>	<b>0.01-25</b>
Leave-On	6	4	0.001-0.05	NR	2	NR	NR	NR	37	128	0.041-0.2	0.05-25
Rinse-Off	15	1	0.0001-0.05	NR	2	NR	NR	NR	4	NR	NR	0.001
Diluted for (Bath) Use	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Eye Area	NR	2	NR	NR	NR	NR	NR	NR	10	NR	0.041	0.05-2
Incidental Ingestion	NR	NR	NR	NR	NR	NR	NR	NR	3	126	NR	0.3-25
Incidental Inhalation-Spray	5 <sup>a</sup>	NR	0.05	NR	1 <sup>b</sup>	NR	NR	NR	7 <sup>a</sup> ; 14 <sup>b</sup>	NR	NR	0.1 <sup>a</sup>
Incidental Inhalation-Powder	NR	NR	NR	NR	1 <sup>b</sup>	NR	NR	NR	14 <sup>b</sup>	NR	0.2 <sup>c</sup>	NR
Dermal Contact	2	3	0.001	NR	4	NR	NR	NR	34	1	0.041-0.2	0.05-2
Deodorant (underarm)	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Hair - Non-Coloring	19	NR	0.001-0.05	NR	NR	NR	NR	NR	4	NR	NR	0.001-0.1
Hair-Coloring	NR	NR	0.0001-0.0075	NR	NR	NR	NR	NR	NR	NR	NR	NR
Nail	NR	NR	NR	NR	NR	NR	NR	NR	NR	1	NR	2
Mucous Membrane	NR	NR	NR	NR	1	NR	NR	NR	3	126	NR	0.3-25
Baby Products	NR	NR	0.001	NR	NR	NR	NR	NR	NR	NR	NR	NR

NR = Not reported

† Because each ingredient may be used in cosmetics with multiple exposure types, the sum of all exposure types may not equal the sum of total uses.

\* Listed with the nomenclature Triticum Aestivum in the VCRP database.

<sup>a</sup>. It is possible these products may be sprays, but it is not specified whether the reported uses are sprays.

<sup>b</sup>. Not specified whether a powder or a spray, so this information is captured for both categories of incidental inhalation.

<sup>c</sup>. It is possible these products may be powders, but it is not specified whether the reported uses are powders.

**Table 6. Ingredients not reported to be in use.**<sup>27,28</sup>

Triticum Aestivum (Wheat) Leaf Extract  
Triticum Aestivum (Wheat) Peptide  
Triticum Monococcum (Wheat) Stem Water  
Triticum Spelta Seed Water  
Triticum Turgidum Durum (Wheat) Seed Extract  
Triticum Vulgare/Aestivum (Wheat) Grain Extract  
Triticum Vulgare (Wheat) Germ Powder  
Triticum Vulgare (Wheat) Straw Water



## REFERENCES

1. Nikitakis J, Kowcz A. wINCI: International Cosmetic Ingredient Dictionary and Handbook. Washington, DC: Personal Care Products Council; 2019. <http://webdictionary.personalcarecouncil.org/jsp/Home.jsp>. Accessed 1/24/2019.
2. Elder RL (ed.). Final Report of the Safety Assessment for Wheat Flour and Wheat Starch. *JEPT*. 1980;4(4):19-32.
3. Andersen FA (ed.). Annual Review of Cosmetic Ingredient Safety Assessments - 2001/2002. *Int J Toxicol*. 2003;22:32-35.
4. Elder RL (ed.). Final Report of the Safety Assessment for Wheat Germ Glycerides and Wheat Gluten. *JEPT*. 1980;4(4):5-17.
5. Burnett CL, Heldreth B, Bergfeld WF, et al. Safety Assessment of Animal- and Plant-Derived Amino Acids as Used in Cosmetics. *Int J Toxicol*. 2014 Nov-Dec;33(Suppl 4):5S-12A.
6. Johnson W, Heldreth B, Bergfeld WF, et al. Safety Assessment of Polysaccharide Gums as Used in Cosmetics. Cosmetic Ingredient Review. Washington, D.C. 2015. Available from CIR.
7. Burnett CL, Fiume MM, Bergfeld WF, et al. Safety Assessment of Plant-Derived Fatty Acid Oils. *Int J Toxicol*. 2017 Nov-Dec;36(Suppl 3):51S-129S.
8. Elder RL (ed.). Final Report of the Safety Assessment for Wheat Germ Oil. *JEPT*. 1980;4(4):33-45.
9. Burnett CL, Bergfeld WF, Belsito DV, et al. Safety Assessment of Hydrolyzed Wheat Protein and Hydrolyzed Wheat Gluten as Used in Cosmetics. *Int J Toxicol*. 2018;37(55S-66S).
10. Thompson T, Grace T. Gluten in cosmetics: Is there a reason for concern? *J Acad Nutr Diet*. 2012;112(9):1316-1323.
11. United States Department of Agriculture (USDA), Agricultural Research Service, National Plant Germplasm System. Germplasm Resources Information Network (GRIN-Taxonomy). <https://npgsweb.ars-grin.gov/gringlobal/taxon/taxonomysimple.aspx>. 2019. Accessed: 05/31/2019.
12. Shewry PR. Wheat. *J Exp Bot*. 2009;60(6):1537-1553.
13. Cardoso RVC, Fernandes A, Heleno SA, et al. Physicochemical characterization and microbiology of wheat and rye flours. *Food Chem*. 2019 May 15;280:123-129.
14. Mankeviciene A, Jablonskyte-Rasce D, Maiksteniene S. Occurrence of mycotoxins in spelt and common wheat grain and their products. *Food Addit Contam Part A*. 2014;31(1):132-138.
15. Kirby EJM. Botany of the Wheat Plant. In: Curtis BC, Rajaram S, Gomez Macpherson H, eds. *Bread Wheat: Improvement and Production*. Rome: Food and Agriculture Organization of the United Nations; 2002.
16. Brandolini A, Hidalgo A. Wheat germ: Not only a by-product. *Int J Food Sci Nutr*. 2012 Mar;63(Suppl 1):71-74.
17. Bar-Sela G, Cohen M, Ben-Arye E, Epelbaum R. The medical use of wheatgrass: Review of the gap between basic and clinical applications. *Mini Rev Med Chem*. 2015;15(12):1002-1010.
18. Anonymous. 2019. Summary Information - Triticum Vulgare (Wheat) Bran Extract. Unpublished data submitted by the Personal Care Products Council on July 8, 2019.
19. Evans DJ, AL P. Chemistry of wheat proteins and the nature of the damaging substances. *Clin Gastroenterol*. 1974;3(1):199-211.
20. Wojakowska A, Perkowski J, Goral T, Stobiecki M. Structural characterization of flavonoid glycosides from leaves of wheat (*Triticum aestivum* L.) using LC/MS/MS profiling of the target compounds. *J Mass Spectrom*. 2013 Mar;48(3):329-339.

21. Zielinski H, Kozłowska H. Antioxidant activity and total phenolics in selected cereal grains and their different morphological fractions. *J Agric Food Chem*. 2000 Jun;48(6):2008-2016.
22. Bahrami N, Yonekura L, Linforth R, et al. Comparison of ambient solvent extraction methods for the analysis of fatty acids in non-starch lipids of flour and starch. *J Sci Food Agric*. 2014 Feb;94(3):415-423.
23. Anonymous. 2019. Summary Information - Aqueous Mixture Containing 3.5%-12% Triticum Vulgare (Wheat) Germ Extract. Unpublished data submitted by the Personal Care Products Council on August 22, 2019.
24. Piergiovanni AR. Extraction and separation of water-soluble proteins from different wheat species by acidic capillary electrophoresis. *J Agric Food Chem*. 2007 May 16;55(10):3850-3856.
25. Perni S, Calzuola I, Gianfranceschi GL, et al. Biochemical and mass spectrometry recognition of phospholipid-peptide complexes in wheat sprouts extract. *J Pept Sci*. 2011 Nov;17(11):744-750.
26. Lucci P, Pacetti D, Calzuola I, et al. Characterization of phospholipid molecular species and peptide molecules in wheat sprout hydroalcoholic extract. *J Agric Food Chem*. 2013 Nov 27;61(47):11453-11459.
27. U.S. Food and Drug Administration Center for Food Safety & Applied Nutrition (CFSAN). Voluntary Cosmetic Registration Program - Frequency of Use of Cosmetic Ingredients. College Park, MD 2020 2020. Obtained under the Freedom of Information Act from CFSAN; requested as "Frequency of Use Data" January 6 2020; received January 13 2020).
28. Personal Care Products Council. 10/2/2017 2017. Concentration of Use by FDA Product Category: Wheat-Derived Ingredients. Unpublished data submitted by Personal Care Products Council.
29. Personal Care Products Council. 2019. Concentration of Use by FDA Product Category - Wheat Additions. Unpublished data submitted by the Personal Care Products Council on October 16, 2019.
30. Johnsen M. The Influence of Particle Size. *Spray Technology and Marketing*. 2004;14(11):24-27.
31. Rothe H. Special Aspects of Cosmetic Spray Evaluation. 2011. Unpublished data presented at the 26 September Expert Panel for Cosmetic Ingredient Safety meeting. Washington, D.C.
32. Bremmer H, Prud'homme de Lodder L, Engelen J. Cosmetics Fact Sheet: To assess the risks for the consumer; Updated version for ConsExpo 4. Bilthoven, Netherlands 2006. Netherlands National Institute for Public Health and the Environment RIVM 320104001/2006. Pages 1-77. <http://www.rivm.nl/bibliotheek/rapporten/320104001.pdf>. Accessed 8/24/2011.
33. Rothe H, Fautz R, Gerber E, et al. Special aspects of cosmetic spray safety evaluations: Principles on inhalation risk assessment. *Toxicol Lett*. 2011;205(2):97-104.
34. CIR Science and Support Committee of the Personal Care Products Council (CIR SSC). 11/3/2015 2015. Cosmetic Powder Exposure. Unpublished data submitted by the Personal Care Products Council.
35. Aylott R, Byrne G, Middleton J, Roberts M. Normal use levels of respirable cosmetic talc: Preliminary study. *Int J Cosmet Sci*. 1976;1(3):177-186.
36. Russell R, Merz R, Sherman W, Siverston J. The determination of respirable particles in talcum powder. *Food Cosmet Toxicol*. 1979;17(2):117-122.
37. European Union. Regulation (EC) No. 1223/2009 of the European Parliament and of the Council of 30 November 2009 on Cosmetic Products. 2009. <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2009:342:0059:0209:en:PDF>. Accessed 11/9/2017.
38. Food Allergen Labeling and Consumer Protection Act of 2004 (FALCPA). 2004. *21 USC 301*. Vol Public Law 108-282, Title II.
39. Sanguigno L, Casamassa A, Funel N, et al. Triticum vulgare extract exerts an anti-inflammatory action in in vitro models of inflammation in microglial cells. *PLoS ONE*. 2018;13(6):e0197493.

40. U.S. Food and Drug Administration (FDA). OTC Active Ingredients. 2010. <https://www.fda.gov/media/75758/download>. Accessed 05/31/2019.
41. Borah M, Sarma P, Das S. A study of the protective effect of *Triticum aestivum* L. in an experimental animal model of chronic fatigue syndrome. *Pharmacognosy Res.* 2014;6(4):285-291.
42. Bonfili L, Amici M, Cecarini V, et al. Wheat sprout extract-induced apoptosis in human cancer cells by proteasomes modulation. *Biochimie.* 2009 Sept;91(9):1131-1144.
43. Telekes A, Hegedus M. Avemar (wheat germ extract) in cancer prevention and treatment. *Nutr Cancer.* 2009;61(6):891-899.
44. Tudek B, Peryt B, Miloszewska J, Szymczyk T, Przybyszewska M, Janik P. The effect of wheat sprout extract on benzo( $\alpha$ )pyrene and 7,2-dimethylbenz( $\alpha$ )anthracene activity. *Neoplasma.* 1988;35(5):515-523.
45. Francois ISJA, Lescroart O, Veraverbeke WS, et al. Safety assessment of a wheat bran extract containing arabinoxylan-oligosaccharides: Mutagenicity, clastogenicity, and 90-day rat-feeding studies. *Int J Toxicol.* 2010;29(5):479-495.
46. European Chemicals Agency (ECHA). Wheat, ext. . <https://echa.europa.eu/registration-dossier/-/registered-dossier/26771/7/7/2> 2019. Accessed: 07/30/2019.
47. Jacobs LR. Enhancement of rat colon carcinogenesis by wheat bran consumption during the stage of 1,2-dimethylhydrazine administration. *Cancer Res.* 1983 Sept;43(9):4057-4061.
48. Arya P, Kumar M. Chemoprevention by *Triticum aestivum* of mouse skin carcinogenesis induced by DMBA and croton oil - association with oxidative status. *Asian Pac J Cancer Prev.* 2011;12(1):143-148.
49. Product Investigations Inc. 2010 2010. Determination of the irritating and sensitizing propensities of a face and body powder (containing 13% *Triticum Vulgare* (Wheat) Germ Extract) on human skin. Unpublished data submitted by Personal Care Products Council.
50. Meding B, Ahonen K, Brisman J, et al. Late skin-prick-test reactions to malted wheat. Clinical observations and immunohistochemical characterization. *Allergy.* 1998 Mar;53(3):282-288.
51. Heederik D, Houba R. An exploratory quantitative risk assessment for high molecular weight sensitizers: Wheat flour. *Ann Occup Hyg.* 2001;45(3):175-185.
52. Prichard MG, Ryan G, Walsh BJ, Musk AW. Skin test and RAST responses to wheat and common allergens and respiratory disease in bakers. *Clin Allergy.* 1985 Mar;15(2):203-210.
53. Brisman J. Baker's Asthma. *Occup Environ Med.* 2002;59(7):498-502.
54. Quirce S, Diaz-perales A. Diagnosis and management of grain-induced asthma. *Allergy Asthma Immunol Res.* 2013;5(6):348-356.
55. Lionetti E, Catassi C. New clues in celiac disease epidemiology, pathogenesis, clinical manifestations, and treatment. *Int Rev Immunol.* 2011;30(4):219-231.
56. Leonard MM, Vasagar B. US perspective on gluten-related diseases. *Clin Exp Gastroenterol.* 2014;7:25-37.
57. University of Chicago Celiac Disease Center. Celiac Disease Facts and Figures. [https://www.cureceliacdisease.org/wp-content/uploads/341\\_CDCFactSheets8\\_FactsFigures.pdf](https://www.cureceliacdisease.org/wp-content/uploads/341_CDCFactSheets8_FactsFigures.pdf). Last Updated Accessed 02/20/2020.
58. Zuidmeer L, Goldhahn K, Rona RJ, et al. The prevalence of plant food allergies: A systemic review. *J Allergy Clin Immunol.* 2008;121(5):1210-1218.
59. Vierk KA, Koehler KM, Fein SB, Street DA. Prevalence of self-reported food allergy in American adults and use of food labels. *J Allergy Clin Immunol.* 2007;119(6):1504-1510.

60. Cianferoni A. Wheat allergy: Diagnosis and management. *J Asthma Allergy*. 2016;9:13-25.
61. Salvatori N, Reccardini F, Convento M, et al. Asthma induced by inhalation of flour in adults with food allergy to wheat. *Clin Exp Allergy*. 2008;38(8):1349-1356.